

Scattering of neutral fermions by a pseudoscalar potential step in two-dimensional space-time

Antonio S. de Castro*
UNESP, Campus de Guaratinguetá
Departamento de Física e Química
Caixa Postal 205
12516-410 Guaratinguetá SP, Brasil

Abstract

The problem of scattering of neutral fermions in two-dimensional space-time is approached with a pseudoscalar potential step in the Dirac equation. Some unexpected aspects of the solutions beyond the absence of Klein's paradox are presented. An apparent paradox concerning the uncertainty principle is solved by introducing the concept of effective Compton wavelength. Added plausibility for the existence of bound-state solutions in a pseudoscalar double-step potential found in a recent Letter is given.

*E-mail address: castro@feg.unesp.br (A.S. de Castro)

In a recent Letter the problem of confinement of neutral fermions by a double-step potential with pseudoscalar coupling in 1+1 dimensions was approached [1]. There it was found that this sort of potential results in confinement in the event that the coupling is strong enough. The one-dimensional Dirac equation developed in that Letter could also be obtained from the four-dimensional one with the mixture of spherically symmetric scalar, vector and anomalous magnetic interactions. If we had limited the fermion to move in the x -direction ($p_y = p_z = 0$) the four-dimensional Dirac equation would decompose into two equivalent two-dimensional equations with 2-component spinors and 2×2 matrices [2]. Then, there would result that the scalar and vector interactions would preserve their Lorentz structures whereas the anomalous magnetic interaction would turn out to be a pseudoscalar interaction. Furthermore, in the (1+1)-world there is no angular momentum so that the spin is absent. Therefore, the (1+1)-dimensional Dirac equation allow us to explore the physical consequences of the negative energy states in a mathematically simpler and more physically transparent way.

In the present Letter some aspects of the scattering of neutral fermions in a pseudoscalar potential step in a (1+1) dimensional space-time are analyzed. This essentially relativistic problem gives rise to some unexpected results and make more credible the existence of the bound-state solutions found in the former Letter [1].

Let us begin by presenting the Dirac equation in 1+1 dimensions. In the presence of a time-independent potential the (1+1)-dimensional time-independent Dirac equation for a fermion of rest mass m reads

$$\mathcal{H}\Psi = E\Psi \quad (1)$$

$$\mathcal{H} = c\alpha p + \beta mc^2 + \mathcal{V} \quad (2)$$

where E is the energy of the fermion, c is the velocity of light and p is the momentum operator. α and β are Hermitian square matrices satisfying the relations $\alpha^2 = \beta^2 = 1$, $\{\alpha, \beta\} = 0$. From the last two relations it steams that both α and β are traceless and have eigenvalues equal to -1 , so that one can conclude that α and β are even-dimensional matrices. One can choose the 2×2 Pauli matrices satisfying the same algebra as α and β , resulting in a 2-component spinor Ψ . The positive definite function $|\Psi|^2 = \Psi^\dagger \Psi$, satisfying a continuity equation, is interpreted as a probability position density and its

norm is a constant of motion. This interpretation is completely satisfactory for single-particle states [3]. We use $\alpha = \sigma_1$ and $\beta = \sigma_3$. For the potential matrix we consider

$$\mathcal{V} = 1V_t + \beta V_s + \alpha V_e + \beta\gamma^5 V_p \quad (3)$$

where 1 stands for the 2×2 identity matrix and $\beta\gamma^5 = \sigma_2$. This is the most general combination of Lorentz structures for the potential matrix because there are only four linearly independent 2×2 matrices. The subscripts for the terms of potential denote their properties under a Lorentz transformation: t and e for the time and space components of the 2-vector potential, s and p for the scalar and pseudoscalar terms, respectively. It is worth to note that the Dirac equation is invariant under $x \rightarrow -x$ if $V_e(x)$ and $V_p(x)$ change sign whereas $V_t(x)$ and $V_s(x)$ remain the same. This is because the parity operator $P = \exp(i\theta)P_0\sigma_3$, where θ is a constant phase and P_0 changes x into $-x$, changes sign of α and $\beta\gamma^5$ but not of 1 and β .

Defining the spinor ψ as

$$\psi = \exp\left(\frac{i}{\hbar}\Lambda\right)\Psi \quad (4)$$

where

$$\Lambda(x) = \int^x dx' \frac{V_e(x')}{c} \quad (5)$$

the space component of the vector potential is gauged away

$$\left(p + \frac{V_e}{c}\right)\Psi = \exp\left(\frac{i}{\hbar}\Lambda\right)p\psi \quad (6)$$

so that the time-independent Dirac equation can be rewritten as follows:

$$H\psi = E\psi \quad (7)$$

$$H = \sigma_1 cp + \sigma_2 V_p + \sigma_3 (mc^2 + V_s) + 1V_t \quad (8)$$

showing that the space component of a vector potential only contributes to change the spinors by a local phase factor.

Provided that the spinor is written in terms of the upper and the lower components

$$\psi = \begin{pmatrix} \phi \\ \chi \end{pmatrix} \quad (9)$$

the Dirac equation decomposes into :

$$\begin{aligned} (V_t - E + V_s + mc^2) \phi(x) &= i\hbar c \chi'(x) + iV_p \chi(x) \\ (V_t - E - V_s - mc^2) \chi(x) &= i\hbar c \phi'(x) - iV_p \phi(x) \end{aligned} \quad (10)$$

where the prime denotes differentiation with respect to x . In terms of ϕ and χ the spinor is normalized as $\int_{-\infty}^{+\infty} dx (|\phi|^2 + |\chi|^2) = 1$, so that ϕ and χ are square integrable functions. It is clear from the pair of coupled first-order differential equations (10) that both $\phi(x)$ and $\chi(x)$ must be discontinuous wherever the potential undergoes an infinite jump and have opposite parities if the Dirac equation is invariant under $x \rightarrow -x$. In the nonrelativistic approximation (potential energies small compared to the rest mass) Eq. (10) becomes

$$\chi = \frac{p}{2mc} \phi \quad (11)$$

$$\left(-\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + V_t + V_s \right) \phi = (E - mc^2) \phi \quad (12)$$

Eq. (11) shows that χ is of order $v/c \ll 1$ relative to ϕ and Eq. (12) shows that ϕ obeys the Schrödinger equation without any contribution from the pseudoscalar potential.

Now, let us choose the nonconserving-parity pseudoscalar potential step

$$V_p(x) = \begin{cases} 0, & x < 0 \\ V_0, & x > 0 \end{cases} \quad (13)$$

Then the solutions of the Dirac equation describing a fermion coming in from the left can be written as

$$\psi(x) = \begin{cases} A \begin{pmatrix} 1 \\ \mu_1 \end{pmatrix} e^{+ik_1 x} + A' \begin{pmatrix} 1 \\ -\mu_1 \end{pmatrix} e^{-ik_1 x}, & x < 0 \\ B \begin{pmatrix} 1 \\ i\mu_2 \end{pmatrix} e^{+ik_2 x}, & x > 0 \end{cases} \quad (14)$$

where

$$\begin{aligned}
(\hbar ck_1)^2 &= E^2 - m^2 c^4 \\
(\hbar ck_2)^2 &= E^2 - m^2 c^4 - V_0^2 \\
\mu_1 &= \frac{\hbar ck_1}{E + mc^2} \\
\mu_2 &= \frac{-i\hbar ck_2 + V_0}{E + mc^2}
\end{aligned} \tag{15}$$

The second line of (15) can be written in the more suggestive form

$$(\hbar ck_2)^2 = E^2 - m^{*2} c^4 \tag{16}$$

where the effective mass

$$m^* = \sqrt{m^2 + V_0^2/c^2} \tag{17}$$

is independent of the sign of V_0 .

The spinor for $x < 0$ describes the incident wave moving to the right (k_1 is a real number) and the reflected wave moving to the left. The spinor for $x > 0$ describes a transmitted wave moving to the right if k_2 is a real number, otherwise it describes an attenuated exponentially wave beyond the potential boundary

The continuity of the spinor at the potential boundary yields the relative amplitudes

$$\begin{aligned}
r &\equiv \frac{A'}{A} = \frac{\hbar c(k_1 - k_2) - iV_0}{\hbar c(k_1 + k_2) + iV_0} \\
t &\equiv \frac{B}{A} = \frac{2\hbar ck_1}{\hbar c(k_1 + k_2) + iV_0}
\end{aligned} \tag{18}$$

In order to determinate the reflection and transmission coefficients we use the probability current density $J = c\psi^\dagger \alpha \psi$, where $\psi^\dagger$ is the Hermitian conjugate spinor. The probability current densities in the two different regions of the x -axis are

$$\begin{aligned}
J(x < 0) &= \frac{2\hbar c^2 k_1}{E + mc^2} (|A|^2 - |A'|^2) \\
J(x > 0) &= \frac{2\hbar c^2 \Re(k_2)}{E + mc^2} |B|^2
\end{aligned} \tag{19}$$

so that the x -independent current density allows us to define the reflection and transmission coefficients as

$$\begin{aligned}
R &\equiv |r|^2 = \frac{\hbar^2 c^2 [k_1^2 + |k_2|^2 - 2k_1 \Re(k_2)] + V_0^2 + 2\hbar c V_0 \Im(k_2)}{\hbar^2 c^2 [k_1^2 + |k_2|^2 + 2k_1 \Re(k_2)] + V_0^2 + 2\hbar c V_0 \Im(k_2)} \\
T &\equiv \frac{\Re(k_2)}{k_1} |t|^2 = \frac{4\hbar^2 c^2 k_1 \Re(k_2)}{\hbar^2 c^2 [k_1^2 + |k_2|^2 + 2k_1 \Re(k_2)] + V_0^2 + 2\hbar c V_0 \Im(k_2)}
\end{aligned} \tag{20}$$

One can note that $R + T = 1$ and R is always less than one, so that Klein's paradox is lacking.

There are two clearly marked behaviors for the spinors. If $E^2 < V_0^2 + m^2 c^4$ then k_2 becomes purely imaginary and $\psi(x > 0)$ describes an evanescent wave ($E < m^* c^2$) with $R = 1$ and $T = 0$. Even though $\psi(x > 0)$ exists there is no probability current density through the potential frontier. Conversely, if $E^2 > V_0^2 + m^2 c^4$ then k_2 becomes a real number so that $\psi(x > 0)$ in fact describes a transmitted wave ($E > m^* c^2$), with $R < 1$ and $T \neq 0$.

Now let us move on to analyze the condition $E^2 > V_0^2 + m^2 c^4$ which ensures that the spinor is an oscillatory wave (k_2 is a real number) in the region $x > 0$. The resulting condition turns into $-\sqrt{E^2 - m^2 c^4} < V_0 < \sqrt{E^2 - m^2 c^4}$ ($m < m^* < E/c^2$). This resembles the nonrelativistic result for a ascending step. On the other side, a deep enough step ($V_0 < -\sqrt{E^2 - m^2 c^4}$) leads to total reflection, in contrast to nonrelativistic quantum mechanics that furnishes $R = 1$ only in the $V_0 \rightarrow -\infty$ case.

The effective mass given by (17) clearly indicates that this kind of potential couples to the mass of the fermion and consequently it couples to the positive-energy component of the spinor in the same way it couples to the negative-energy component. The energy spectrum for a free fermion ranges continuously from $-mc^2$ to $-\infty$ as well as from mc^2 to ∞ and the potential step contributes to enhance the energy gap bringing into being at $x = 0$ an

ascending step for the positive-energy solutions and a descending step for the negative-energy solutions. The spectral gap always stretches whichever the sign of V_0 . This is so because the effective mass depends on V_0^2 . In other words, the potential step at $x = 0$ is repulsive for the particles as well as for the antiparticles. In this way one can see that there is no room for transitions from positive- to negative-energy solutions. It all means that Klein's paradox does not come to the scenario.

Another interesting way of visualizing the absence of Klein's paradox is to consider the uncertainty in the position beyond the potential boundary in the case $E^2 < V_0^2 + m^2 c^4$ ($E < m^* c^2$). In this circumstance the probability density for $x > 0$ is given by

$$\rho(x > 0) = |B|^2 e^{-2|k_2|x} \left[1 + \left(\frac{\hbar c |k_2| + V_0}{E + m c^2} \right)^2 \right] \quad (21)$$

so that

$$\Delta x = \frac{\hbar c}{2\sqrt{m^2 c^4 + V_0^2 - E^2}} \quad (22)$$

From this last result one can see that the penetration of the fermion into the region $x > 0$ will shrink without limit as $|V_0|$ increases and there is no atmosphere for the production of particle-antiparticle pairs at the potential frontier. It means that the localization of a fermion under the influence of this sort of pseudoscalar potential does not require any minimum value in order to ensure the single-particle interpretation of the Dirac equation. At first glance it seems that the uncertainty principle dies away provided such a principle implies that it is impossible to localize a particle into a region of space less than half of its Compton wavelength (see, *e.g.*, Ref [2]). This apparent contradiction can be remedied by writing (22) into the form

$$\Delta x = \frac{\hbar c}{2\sqrt{m^{*2} c^4 - E^2}} \quad (23)$$

Thus, it can be immediately seen that the minimum uncertainty is

$$\Delta x_{\min} = \frac{\hbar}{2m^* c} \quad (24)$$

in the limit as $|V_0| \rightarrow \infty$, and again the sign of V_0 plays no part. Therefore, for obtaining a result consistent with the uncertainty principle it is a must

to use the effective Compton wavelength.

In addition to their intrinsic importance, the above conclusions for the scattering of a neutral fermion by a pseudoscalar single-step potential enhance the plausibility arguments for the confinement of neutral fermions by a pseudoscalar double-step potential presented in the former Letter [1].

We have made the analysis of the scattering of neutral fermions by a pseudoscalar potential step in the (1+1)-dimensional Dirac equation. The results obtained in the (1+1) world are also supposed to be qualitatively true in the more realistic 3+1 world. Although the idea and calculations presented in this Letter are simple the results and conclusions seem to be new.

Acknowledgments

This work was supported in part through funds provided by CNPq and FAPESP.

References

- [1] A.S. de Castro and W.G. Pereira, Phys. Lett. A, to be published.
- [2] P. Strange, Relativistic Quantum Mechanics, Cambridge University Press, Cambridge, 1998.
- [3] B. Thaller, The Dirac Equation, Springer-Verlag, Berlin, 1992.